Lab Use Only				
Res Lab #:				
Date Rec'd:				
Initials:				

ACD (yellow top) or

DNA

National Inherited Bleeding Disorder Genotyping Laboratory

Department of Pathology and Molecular Medicine Queen's University, Kingston, Ontario



von Willebrand Disease Genotype Testing Requisition

Patient Name:	Surname, First Name)	Se	x: Male □ Female	Male □ Female □	
(\$	Surname, First Name)				
DOB://	Unique Iden	tifier:	CBDR#:		
YY MM	DD	eg. Health card #, H	ospital #		
Date of specimen	collection:/_	/ Phlebotomis	t:		
Referring Clinic:	R	Report to:	Fax #: _		
Von Willebrand D	isease: Type 2(subtype if kr		☐ Type 1C		
Testing: VWF:Ag	IU/mL				
	o	☐ Other	_ Value	IU/mL	
Factor V	'III:C IU/mL	-			
Multime	rs				
Desmop	ressin trial				
Pregnant?	Yes □ No □				
Have samples from	n this family been sent to	this lab before? Y	′es □ No □		
If Yes, specify					
Relationship to this	s patient				
Sample Requ	uirements:		Ship to:		
6 cc whole	e blood	Attn: Gina Jones/Samira Kheitan Department of Pathology and Molecular Medicine			
EDTA (laven	der top) <u>or</u>				
		Queen's Univer	rsity, Richardson Labo	ratory, Room 201	

88 Stuart St., Kingston, Ontario K7L 3N6 Tel: 613-533-3187 FAX: 343-344-2733

Email: NIBDGL@queensu.ca

National Inherited Bleeding Disorder Genotyping Lab Sample Collection Instructions

Requisitions:

- 1. Samples must be accompanied by a completed requisition form.
- 2. Submitted patients must have a documented rationale for testing:
 - For affected hemophilia A or B patients, a clotting factor activity level must be provided.
 - For VWD patients, VWF:Ag, VWF activity (specify test used), FVIII:C, and multimer information must be provided.
 - For carrier testing, a documented family history or a coagulation factor level must be provided.
 - For carrier testing, information on the family variant if available must be provided.
 - For prenatal testing, information on the family variant if available must be provided.
- 3. Incomplete requisition forms will result in delayed sample testing.

Sample Collection and Shipment:

- 1. Samples acceptable for testing:
 - Venous whole blood (minimum 6 cc) collected into EDTA (lavender top) or ACD (yellow top) evacuated tubes
 - Expired tubes should not be used.
 - If blood is being drawn from an intravenous line for laboratory testing, two times the deadspace volume should be discarded.
 - When drawing blood specimens for several examinations during a single venipuncture, the "order of draw" shall be: (1) blood culture; (2) coagulation specimens; (3) serum tube with or without clot activator or gel; (4) heparin; (5) EDTA; (6) glycolytic inhibitor.
 - DNA (minimum of 15 μg at 150 ng/μL); for some cases smaller samples are acceptable.
 - Patient ID must be verified and samples labelled using two unique identifiers.
 - Samples shall be collected using routine practices/standard precautions.
 - Materials for sample collection shall be safely disposed of according to institutional protocols.
- 2. Ship packages on a Monday, Tuesday, Wednesday, Thursday as follows:
 - Shipping temperature: DNA (ambient), whole blood (cold packs), whole blood frozen (dry ice).
 - Place the samples in sealable plastic bags with absorbent material.
 - Include completed requisition and consent form.
 - Ship overnight. Contact courier for complete shipping instructions.
- 3. Attach the following labels to the outside of the box/package:
 - Dry ice label (if applicable).
 - Return address label (including the contact name and telephone number).
- 4. Ship to: Attn: Gina Jones/Samira Kheitan

Department of Pathology and Molecular Medicine Queen's University, Richardson Laboratory, Room 201 88 Stuart Street, Kingston, Ontario K7L 3N6

Tel: 613-533-3187 FAX: 343-344-2733 NIBDGL@queensu.ca

Results:

- Turnaround time is approximately 3 months from the time of sample submission but may be longer for rare genes or for carrier testing if familial variant is unknown.
- If the family-specific variant is known, urgent reporting can be completed within several weeks. Please indicate this on the requisition form.
- For prenatal testing, maternal cell contamination studies are the responsibility of the referring clinic.
- Results will be faxed to the referring clinician or designate listed on the requisition.
- Results will also be entered into the Canadian Bleeding Disorders Registry (CBDR) when possible.